


# Chapter #14 – Chemical Kinetics

1. Use the following table to calculate the rate of disappearance for  $X_2$  at different times.



| Time (s) | $[X_2]$ |
|----------|---------|
| 0.00 s   | 1.30 M  |
| 15.0 s   | 1.10 M  |
| 30.0 s   | 0.80 M  |
| 60.0 s   | 0.20 M  |
| 120.0 s  | 0.050 M |

This is specifically disappearance

$$\text{Rate} = \frac{-\Delta [\text{Reactants}]}{\Delta \text{time}}$$

, where  $\Delta$  means  
Final - initial of that  
quantity

Do NOT get  
this confused  
with  
 $\text{rate} = k [\text{stuff}] \dots$

2 SF(s)

a. 0 s  $\rightarrow$  120 s

$$\text{Rate} = \frac{-[(0.050\text{M}) - (1.30\text{M})]}{(120\text{s}) - (0.00\text{s})} = 0.012 \frac{\text{M}}{\text{s}} \quad \text{Molarity per second}$$

b. 0 s  $\rightarrow$  60 s

$$\text{Rate} = \frac{-[(0.20\text{M}) - (1.30\text{M})]}{(60.0\text{s}) - (0.00\text{s})} = 0.018 \frac{\text{M}}{\text{s}}$$

c. 15 s  $\rightarrow$  60 s

$$\text{Rate} = \frac{-[(0.20\text{M}) - (1.10\text{M})]}{(60.0\text{s}) - (15.0\text{s})} = 0.020 \frac{\text{M}}{\text{s}}$$

d. What would the rate of disappearance be for Y at these time intervals?

The rate of disappearance for Y would be  $\frac{2}{3}X$  the rate of disappearance for  $X_2$ .



2. Give the  $t_{1/2}$  equation. Half Life, scary.

$$t_{1/2} = \frac{\ln(2)}{K} \quad \text{or} \quad t_{1/2} = \frac{0.693}{K}$$

Where K is  
the rate  
constant.

3. What effect does temperature have on the rate constant (k)?

Temperature and the magnitude of the  
rate constant, K, are DIRECTLY  
Proportional.



$$\frac{2X}{3} = \frac{3Y}{3}$$


$$Y = \frac{2}{3}X$$



4. How does the activation energy affect the rate constant (k)?

A high activation energy results in a low rate constant; it takes MORE energy for the rxn to proceed.

5. Label the following graph:



The calculus  
is integration

1<sup>st</sup> order reaction: natural log order

$$\text{rate} = k[A] \rightarrow \text{Calculus} \rightarrow \ln[A]_t = \ln[A]_0 - kt$$

from this, a plot of  $\ln[A]$  vs. time makes a straight line if the rxn is 1<sup>st</sup> order.


2<sup>nd</sup> order reaction: inverse order

$$\text{rate} = k[A]^2 \rightarrow \text{Calculus} \rightarrow \frac{1}{[A]_t} = kt + \frac{1}{[A]_0}$$

from this, a plot of  $\frac{1}{[A]}$  vs. time makes a straight line if the rxn is 2<sup>nd</sup> order.



1. Consider the data for hydrogen concentration  $[H_2]$ , iodine concentration  $[I_2]$  and the rate of reaction (moles per liter per second or  $\text{mol}\cdot\text{L}^{-1}/\text{s}$ ) for this reaction:



| Trial | $[H_2]$ (mol/L) | $[I_2]$ (mol/L) | Rate ( $\text{mol}\cdot\text{L}^{-1}/\text{s}$ ) |
|-------|-----------------|-----------------|--------------------------------------------------|
| 1     | 0.01            | 0.05            | 0.04                                             |
| 2     | 0.02            | 0.05            | 0.08                                             |
| 3     | 0.05            | 0.01            | 0.02                                             |
| 4     | 0.05            | 0.03            | 0.54                                             |

$$2^n = 2$$

$$n = 1$$

So,  $[H_2]^1$

$$3^n = 27$$

$$n = 3$$

So,  $[I_2]^3$


What is the overall reaction order?

$$\text{rate} = k[\text{stuff}] \dots$$

$$\text{rate} = k[H_2]^1[I_2]^3, \text{ add 1 and 3;}$$

The overall order is 4.

2. Consider a hypothetical reaction:



- ① Doubling the concentration of A causes the reaction rate to increase by a factor of four. This is done while the concentration of B is held constant. Tripling the concentration of B, while the concentration of A is held constant, causes the reaction rate to increase by a factor of nine. What is the rate law expression for this reaction?

②

$$2A = 4\text{rate} ; 3B = 9\text{rate}$$

$$2n = 4$$

$$n = 2$$


$$3n = 9$$

$$n = 2$$

$$\left. \begin{array}{l} 2A = 4\text{rate} \\ 3B = 9\text{rate} \end{array} \right\} \text{rate} = k[A]^2[B]^2$$



3. Consider the reaction:



| Trial | $[\text{NO}_2^-] (\text{mol/L})$ | $[\text{NH}_4^+] (\text{mol/L})$ | Rate ( $\text{mol} \cdot \text{L}^{-1} / \text{s} \times 10^{-7}$ ) |
|-------|----------------------------------|----------------------------------|---------------------------------------------------------------------|
| 1     | 0.0100                           | 0.200                            | 5.4                                                                 |
| 2     | 0.0200                           | 0.200                            | 10.8                                                                |
| 3     | 0.0400                           | 0.200                            | 21.6                                                                |
| 4     | 0.200                            | 0.200                            | 10.8                                                                |
| 5     | 0.200                            | 0.400                            | 21.6                                                                |
| 6     | 0.200                            | 0.600                            | 32.4                                                                |

$$2n = 2$$

$$n = 1$$

$$2n = 2$$

$$n = 1$$

Determine the rate law expression for this reaction:

So,  $\text{rate} = k [\text{NO}_2^- (\text{aq})] [\text{NH}_4^+ (\text{aq})]$

Use trial #1:

$$\frac{(5.4 \frac{\text{M}}{\text{s}})}{(0.0100 \text{M})(0.200 \text{M})} = \frac{k(0.0100 \text{M})(0.200 \text{M})}{(0.0100 \text{M})(0.200 \text{M})}$$

4. In a reaction involving only one reactant, A, the rate of the reaction increases by a factor of 27 when the concentration of A is tripled. What is the rate law expression for this reaction?